Medication Safety in Older Adults: Excellent Care for Geriatric Patients in the Community Setting

Michelle A Fritsch, PharmD, BCGP
Representing the Peter Lamy Center on Drug Therapy and Aging
Founder Retirement Wellness Strategies
Co-Founder Propel Comprehensive Wellness
Co-Founder Medipreneurs
Harrisburg, NC

Objectives

- 1. Describe the impact of pharmacist intervention over 3 decades of a patient case.
- 2. Identify three ways to take a preventive vs reactive approach to healthy aging with plans to incorporate these in your practice.
- 3. Incorporate the team that can help you achieve your clinical goals and help your patient meet therapeutic goals.



How many have at least 25% geriatric patients in practice?

Yes = raise your hand high


No = keep your hand down

Some interesting facts

- Over 65 average TV consumption 7.25 hours/day
- 10,000 turning 65 each day







How many have a falls risk reduction program?

Comprehensive program = raise both hands high

Some falls risk reduction = raise one hand high

Nothing for fall yet, but open to ideas = wave your hand in the air

I don't see that happening = keep your hand down

How many have a patient's full medication list in your system?

Rx, OTC, supplements, & anything else updated= raise both hands high

Rx and OTC = raise one hand high

Just Rx, but open to ideas = wave your hand in the air

Rx is all we keep = keep your hand down

prescription

Over-the-counter

Tobacco

Vitamins

Do you know ALL medicines/substances your patients take?

caffeine

Herbal

Supplements


Marijuana | CBD

Holistic

Drug Interactions

- <u>Drug interactions</u> have been shown to cause a <u>decline in functional</u> <u>abilities</u> in older people. Drugs can interact to alter the absorption, distribution, metabolism, or excretion of a drug or interact in a synergistic or antagonist fashion altering their pharmacodynamics.
- Increase ER visits
- Increase hospitalizations
- Increase physician visits

Crit Rev Oncol Hematol. 2003 Nov;48(2):133-43.



How well do you really know your patients?

Cost

At risk of isolation?

Understand directions Remember to take it

Who is backup?

Safety of environment

Able to open the bottle

Able to get to the pharmacy

Providing more Whole-Person Geriatric Care

'We draw from decades of accumulated knowledge, skills, and experience in areas that are central to geriatric medicine, including expertise in complexity and the biopsychosocial model; attention to function and quality of life; the ability to provide culturally competent, person-centered care; the ability to assess people's preferences and values; and understanding the importance of systems in optimizing outcomes.' J Am Geriatr Soc 67:17–20, 2019.

Lee is an 82-year-old living in assisted living; ambulates w/rollator

Past Medical History

- Severe knee & shoulder osteoarthritis, s/p arthroplasty on right knee
- s/p left hip arthroplasty after a fall when left knee gave out
- Urinary incontinence
- Hypertension
- Type 2 diabetes
- Depression
- BMI 32

Current Medications

- Oxycodone 5mg every 6 hours
- Naproxen 220mg two every 12 hours
- Gabapentin 300mg three times daily
- Oxybutinin ER 15mg daily
- Losartan 100mg daily
- Metoprolol succinate 100mg daily
- Metformin XR 2000mg daily
- Glimepiride 1mg daily
- Sertraline 50mg daily
- Medical cannabis 14:1 gummies 2 twice daily

Let's discuss.

What are your concerns?

Where would you begin?

- Falls
- Falls Risk Inducing Drugs
- Falls risk associated medical conditions
- Bone health

- Risk for isolation
- Quality of life
- Purpose*

Lee is an 82 /ear-old living in assisted living; ambulates w/ rollator

Past Medical History

- Severe knee & shoulder osteoarthritis, s/p arthroplasty on right knee
- s/p left hip arthroplasty after a fall when left knee gave out
- Urinary incontinence Functional? Stress? Prostate?

BP? A1c?

- Hypertension
- Type 2 diabetes
- Depression
- BMI 32 <

Weight changes?

Current Medications

- Oxycodone 5mg every 6 hours
- Naproxen 220mg two every 12 hours
- Gabapentin 300mg three times daily
- Oxybutinin ER 15mg daily
- · Losartan 100mg daily
- Metoprolol succinate 100mg daily
- Metformin XR 2000mg daily
- Glimepiride 1mg daily
- Sertraline 50mg daily
- Medical cannabis 14:1 gummies 2 twice daily

If you were performing a review in this assisted living facility, how much time would you spend on this review?

Lee is a 72-year-old living at home

Past Medical History

- Moderate knee & shoulder osteoarthritis, planned arthroplasty on right knee
- Urinary incontinence
- Hypertension
- BMI 28.8

Current Medications

- Naproxen 220mg two every 12 hours
- Gabapentin 300mg twice daily
- Oxybutinin ER 15mg daily
- Lisinopril 10mg daily
- Metoprolol succinate 25mg daily

Let's discuss.

What are your concerns?

Where would you begin?

- Falls risks?
- FRIDs?
- Falls risk with medical conditions?
- Bone health?

- Risk for isolation
- Quality of life
- Exercise, Diet
- Social health
- Purpose*

Lee is 72 ear-old living at home

Past Medical History

- Moderate knee & shoulder osteoarthritis, planned arthroplasty on right knee
- Urinary incontinence Functional? Stress? Prostate?
- Hypertension ← BP?
- BMI 28.8 Weight changes?

Current Medications

- Naproxen 220mg two every 12 hours
- Gabapentin 300mg twice daily
- Oxybutinin ER 15mg daily
- Lisinopril 10mg daily
- Metoprolol succinate 25mg daily

What if an opioid was added?

Lee is a 62-year-old living at home

Past Medical History

- Mild knee & shoulder osteoarthritis
- Hypertension
- BMI 24.8

Current Medications

- Naproxen 220mg one as needed
- Lisinopril 10mg daily


Let's discuss.

What are your concerns?


Where would you begin?

- Falls Risks?
- Bone Health?

- Risk for isolation
- Quality of life
- Exercise, Diet
- Social health
- Purpose*



If we provide more wholeperson geriatric care, what does that entail?



Asking more questions

Thinking more about prevention

When and how do we best incorporate pharmacogenomics and other precision medicine elements?

What non-medication options could be added to the therapeutic plan? What is our role as pharmacists?

Where do we fit in integrative medicine?

How can we help prevent falls?

How can we help prevent functional decline?

How can we help prevent weight gain?

Creating a solid team

Local primary care providers

Local geriatricians, or within the state

Physical therapists

Occupational therapists

Dieticians



Knowing local resources

- Aging Lifecare Managers https://www.aginglifecare.org/
- Professional Senior Move Managers <u>https://www.nasmm.org/</u>
- Certified aging in place (CAPS) home modification experts <u>https://www.nahb.org/education-and-events/education/designations/Certified-Aging-in-Place-Specialist-CAPS</u>
- Places to keep pets short or long term
- Safe places to exercise
- Senior specialist transportation options
- Area Agency on Aging, senior centers

Medication Safety Includes...

- Drug interactions with other meds, food, or conditions
- Allergies
- Untreated medical indication
- Not the best medication choice for this patients
- Subtherapeutic dose
- Excessive dose
- Failure to receive the medications
- Medication without an indication

- Inappropriate prescribing
- Inappropriate delivery
- Inappropriate patient behavior
- Patient idiosyncrasy
- Inappropriate monitoring

Lee is a 68-year-old living at home

Past Medical History

- Mild knee & shoulder osteoarthritis
- Hypertension
- BMI 24.8

Current Medications

- Naproxen 220mg one as needed
- Lisinopril 10mg daily



In case you're interested... www.rapidalliance.org

